

Mossbauer study of cation substitution effect on the magnetic phase transition temperature of 0.5AFeO₃-0.5NaNbO₃ (A = Bi, La) solid solutions

S.I. Raevskaya¹, S.P. Kubrin¹, Jian Zhuang², I.P. Raevski¹,
M.A. Malitskaya¹, I.N. Zakharchenko¹, V.V. Titov¹, V.G. Smotrakov¹

¹*Research Institute of Physics and Faculty of Physics, Southern Federal University,
344090, Rostov-on-Don, Russia
sveta.raevskaya@mail.ru*

²*Electronic Materials Research Laboratory, Key Laboratory of the Ministry of Education and
International Center for Dielectric Research, Xi'an Jiaotong University, 710049, Xi'an, P.R. China*

BiFeO₃ is the most widely studied multiferroic due to high temperatures of both ferroelectric and magnetic phase transitions. Recently a possibility of magnetic superexchange between Fe³⁺ ions in BiFeO₃ via the empty 6p states of Bi³⁺ was predicted theoretically [1]. Earlier we showed the role of Pb²⁺ ions which also possess the empty 6p states, in the enhancement of the T_N in perovskite Pb_{1-x}A_xFe_{1/2}Nb_{1/2}O₃ (A = Ca, Ba) solid solutions [2, 3]. However this mechanism of superexchange seems to become noticeable only at high enough dilution of the Fe-sublattice, i.e. in solid solutions.

The 0.5AFeO₃-0.5NaNbO₃ (A = Bi, La) solid solution compositions were prepared by solid phase reactions route from high-purity oxides. The XRD study revealed that the samples are single phase and perovskite structure. Room temperature Mossbauer spectra of both show the quadrupole-split lines. The main reason of quadrupole splitting in such systems is compositional disorder in B-sublattice. All spectra contain 2 or 3 doublets with different values of quadrupole splitting. The isomer shift values of doublets correspond to the Fe³⁺ ions in oxygen octahedron. The presence of 2 or 3 doublets indicates that Fe³⁺ has 2 or 3 different environments in the lattice which may be a fingerprint of B-cations short-range ordering or clusterization [4].

To estimate the temperature of magnetic phase transition (T_N) we performed the measurement of Mossbauer spectrum line intensity under subsequent temperature lowering. Near magnetic phase transition the Mossbauer spectrum transforms from doublet into sextet. This transformation is accompanied by a dramatic decrease of Mossbauer spectrum line intensity I_m . Thus, the abrupt drop in the $I_m(T)$ dependence corresponds to the T_N . Mossbauer studies have shown that while for 0.5BiFeO₃-0.5NaNbO₃ the T_N value is about 150 K, for 0.5LaFeO₃-0.5NaNbO₃ it is only ≈ 20 K. This dramatic difference in T_N values seems to be due to additional contribution of the superexchange between Fe³⁺ ions via the empty 6p states of Bi³⁺ to the overall superexchange in accord with predictions of De Sousa et al. [1].

For both compositions magnetization M studies in the 5-200 K range revealed only a maximum at 20-25 K in the zero-field cooling mode. In the field-cooled mode this maximum was not observed. Such behavior is typical of spin glasses. In PbFe³⁺_{1/2}B⁵⁺_{1/2}O₃ (B⁵⁺ = Nb, Ta) perovskite multiferroics a spin-glass state is known to coexist at low temperatures with the antiferromagnetic state [2, 4]. The absence of the $M(T)$ anomaly at around 150 K for 0.5BiFeO₃-0.5NaNbO₃ composition seems to be due to the strong diffusion of this anomaly. Similar "disappearance" of the $M(T)$ anomaly corresponding to antiferromagnetic phase transition was observed, e.g. for some compositions of the PbFe_{2/3}W_{1/3}O₃-PbTiO₃ and PbFe_{1/2}Nb_{1/2}O₃-PbTiO₃ solid solutions [5, 6].

This work was supported in part by RFBR project 19-52-53030 GFEN_a and by the Ministry of Education and Science of the Russian Federation (research projects 3.1649.2017/4.6 and 3.5346.2017/BP).

1. R. De Sousa, M. Allen, M. Cazayous, *Phys. Rev. Lett.* **110**, 267202 (2013).
2. I.P. Raevski, S.P. Kubrin, S.I. Raevskaya, et. al., *Phys. Rev. B*, **80** (2009).
3. I.P. Raevski, S.P. Kubrin, S.I. Raevskaya, et. al., *Ferroelectrics* **398** (2010).
4. I.P. Raevski, S.P. Kubrin, S.I. Raevskaya, et. al., *Phys. Rev. B* **85** (2012).
5. L. Mitoseriu, D. Marre, A. S. Siri, P. Nanni, *Appl. Phys. Lett.* **83**, 5509 (2003).
6. S.P. Singh, S.M. Yusuf, S. Yoon, et. al., *Acta Mater.* **58**, 5381 (2010).